

REF.	No.	SPECIES	AUTHOR	GRADE	Ø MIN.(mm)	Ø MAX.(mm)	PLACE
F12	2	<i>Brotia costula varicosa</i>	Troschel,1837	F+	50,0	56,0	F
F11	8	<i>Cassidula nucleus</i>	Gmelin,1791	F++/G	18,0	22,0	F
F15		<i>Corbicula fluminea</i>	Müller,1774	F/F+/F++	15,0	25,0	A
F27	1	<i>Cyrenobatissa subsulcata</i>	Clessin,1878	F+ (usual)		49,0	CH
F9	1	<i>Faunus ater</i>	Born,1758	F++		62,0	F
F24		<i>Galba truncatula</i>	Müller,1774	F+/F++	2,0	5,0	J
F22		<i>Hydrobia ulvae</i>	Pennant,1777	G	2,0	10,0	LC/H
F21	8	<i>Melanoides tuberculata</i>	Müller,1774	F++	11,0	25,0	RD
F13	1	<i>Neocorbicula limosa</i>	Maton,1809	F++		15,0	UR
F20	2	<i>Ovatella myosotis</i>	Draparnaud,1801	F++		6,0	LC
F26		<i>Physella acuta</i>	Draparnaud,1805	F++/G	9,0	14,0	LC
F17	6	<i>Planorbis planorbis</i>	Linné,1758	F+	2,0	5,0	BA
F10	7	<i>Sadleriana pannonica</i>	Frauenfeld,1865	F++		1,0	ES
G356	1	<i>Solenaia carinata</i>	Heude,1877	F+ (usual)		224,0	CH
F19	1	<i>Unio douglassiae taiwanicus</i>	Pilsbry,1921	F+		72,0	CH

Véase identificadores de localidades en la sección "Leyenda por países"

See place identifiers in the "Countries Caption" section