

REF.	No.	SPECIES	AUTHOR	Ø MIN. (mm)	Ø MAX. (mm)	PLACE	VALUE(€)
L14	1	<i>Amphidromus quadrasi quadrasi</i>	Hidalgo,1887		41,0	F	4
L51	1	<i>Balea biplicata</i>	Montagu,1803		19,0	B	2
L40	4	<i>Bradybaena similaris</i>	Ferussac,1821	16,0	17,0	CH	2
L69	1	<i>Bulimulus</i> sp.			17,0	A	8
L16	5	<i>Calocochlia cailliaudi</i>	Deshayes,1839	35,0	45,0	F	4
L47		<i>Candidula gigaxii</i>	Pfeiffer,1850	6,0	11,0	J	2
L49		<i>Caracollina lenticula*</i>	Michaud,1831	5,0	8,0	J/G	1
L41	1	<i>Cathaica fasciola</i>	Draparnaud,1801		16,0	CH	3
L80		<i>Cecilioides acicula</i>	Müller,1774	3,0	5,0	J	1
L33		<i>Cepaea nemoralis</i>	Linné,1758	20,0	28,0	CN/D	1,5
L15		<i>Cerneuella virgata*</i>	Da Costa,1778	8,0	14,0	J/Z	0,5
L73		<i>Chondrina farinesi farinesi</i>	Des Moulins,1835	4,0	7,0	J	2
L79		<i>Chondrina kobelti kobelti</i>	Westerlund,1887	4,0	5,0	CN	1,5
L29	3	<i>Chrysalis virgata cerina</i>	Bartsch,1932	42,0	51,0	F	3,5
L14	1	<i>Chrysalis virgata porracea</i>	Jay,1839		47,0	F	2,5
L52		<i>Cochlicella acuta</i>	Müller,1774	12,0	16,0	J/CN	1
L77		<i>Cochlostoma oscitans</i>	Gofas,1989	10,0	11,0	CN	2,5
L40	1	<i>Cyclophorus batanicus</i>	Quadras & Möllendorff,1894		20,0	F	2
L17		<i>Cyclophorus linguiferus</i>	Sowerby,1843	29,0	34,0	F	2,5
L78	1	<i>Cyclophorus martensianus</i>	Moellendorff,1874		25,0	CH	4
L82	2	<i>Cyclotus caroli</i>	Kobelt,1886	17,0	19,0	F	5
L37	1	<i>Cyclotus taivanus</i> W/O	Adams,1870		11,0	TI	6
L71	3	<i>Discus rotundatus</i>	Müller,1774		5,0	LC	1
C11		<i>Dupotetia tigris albino form *</i>	Servain,1880	20,0	32,0	MC	2,5
C9		<i>Eobania vermiculata</i>	Müller,1774	25,0	29,0	MU	1
L48	1	<i>Euphaedusa aculus</i>	Benson,1842		13,0	CH	3
L53		<i>Ferussacia folliculus*</i>	Linné,1758	6,0	10,0	J	0,5
L44	1	<i>Granaria frumentum</i>	Draparnaud,1801		7,5	HN	2,5
L78		<i>Hatumia cobosi</i>	Ortiz de Zárate,1962	7,0	10,0	AL	8
L85		<i>Helicella stiparum</i>	Rossmässler,1854	13,0	18,0	AL	5
L83		<i>Helicella zujarensis</i>	Ortiz de Zárate,1950	6,0	7,0	AL	8
L13		<i>Helicostyla annulata</i>	Sowerby,1831	19,0	24,0	F	1,5
L9		<i>Helix aspersa</i>	Müller,1774	28,0	36,0	J	0,5
L45	1	<i>Hemiphaedusa thaleroptyx</i>	Moellendorff,1882		19,0	CH	6
L74	2	<i>Iberus alcarazanus</i>	García S. Nicolás,1957	25,0	26,0	J	20
L4		<i>Iberus alonensis white</i>	Ferussac,1821	32,0	40,0	J	2,5
L4		<i>Iberus alonensis cream</i>	Ferussac,1821	32,0	40,0	J/AL	3,5
L86	4	<i>Iberus alonensis subfossil</i>	Ferussac,1821	36,0	38,0	J	4,5
L122		<i>Iberus angustatus type 1</i>	Rossmässler,1854	20,0	27,0	J	5
L121		<i>Iberus angustatus type 2</i>	Rossmässler,1844	20,0	27,0	J	4
L90	3	<i>Iberus campesinus subfossil</i>	Exquerra,1845	35,0	39,0	AL	8
L77	8	<i>Iberus cf. cobosi</i>	Ibáñez & Alonso,1977	19,0	24,0	M	25
L28		<i>Iberus cf. loxanus (costulated)</i>	Rossmässler,1854	22,0	28,0	M	15
L31		<i>Iberus cf. marmoratus type 1</i>	Ferussac,1821	20,0	25,0	M	20
L73	8	<i>Iberus cf. marmoratus type 2</i>	Ferussac,1821	20,0	22,0	M	20
L7		<i>Iberus cobosi type 1</i>	Ibáñez & Alonso,1977	21,0	26,0	M	15
L7	6	<i>Iberus cobosi type 2</i>	Ibáñez & Alonso,1978	21,0	26,0	M	18
L1		<i>Iberus gualtieranus typical form</i>	Linné,1758	17,0	40,0	J	4
L2		<i>Iberus gualtieranus rare forms</i>	Linné,1758	25,0	43,0	J	12
L69	8	<i>Iberus gualtieranus (white and cream)</i>	Linné,1758	36,0	43,0	AL	6
L43	1	<i>Iberus gualtieranus "freak"</i>	Linné,1758		38,0	J	12
L79	4	<i>Iberus gualtieranus subfossil</i>	Linné,1758	31,0	41,0	J	8
L5	5	<i>Iberus gualtieranus subfossil</i>	Linné,1758	39	49	AL	4
L2		<i>Iberus gualtieroloxanus</i>	García San Nicolás,1949	22,0	26,0	M	15
L87		<i>Iberus guiraoanus type 1</i>	Rossmässler,1854	21,0	30,0	J	7
L87		<i>Iberus guiraoanus type 2</i>	Rossmässler,1854	21,0	30,0	J	8
L74	1	<i>Iberus guiraoanus "freak"</i>	Rossmässler,1854		22,0	J	18

L89		<i>Iberus intermedius</i>	Boettger, 1913	29,0	38,0	J	20
L38	4	<i>Iberus intermedius</i> subfossil	Boettger, 1913	32,0	36,0	J	22
L2		<i>Iberus laurenti</i>	Bourguignat, 1870	33,0	39,0	J	15
L124		<i>Iberus loxanus</i> type 1	Rossmässler, 1854	21,0	26,0	G	8
L124		<i>Iberus loxanus</i> type 2	Rossmässler, 1854	19,0	26,0	G	8
L8		<i>Iberus marmoratus</i> type 1	Ferussac, 1821	18,0	25,0	G	4
L8		<i>Iberus marmoratus</i> type 2	Ferussac, 1821	18,0	25,0	G	5
L123		<i>Iberus marmoratus</i> type 3	Ferussac, 1821	19,0	22,0	M	4
L77	5	<i>Iberus marmoratus</i> type 4	Ferussac, 1821	21,0	23,0	M	5
L80	1	<i>Iberus marmoratus</i> "freak"	Ferussac, 1821		21,0	G	7
L32		<i>Iberus minor</i>	Serradeli	32,0	40,0	J	15
L89	8	<i>Iberus minor</i> (white and cream)	Serradeli	39,0	42,0	AL	15
L91		<i>Iberus ornatissimus</i> clear	Cobos, 1979	27,0	32,0	AL	20
L91		<i>Iberus ornatissimus</i> dark	Cobos, 1979	27,0	32,0	AL	25
L10		<i>Iberus ortizi</i> type 1	García San Nicolás, 1957	19,0	28,0	J/CO	7
L10		<i>Iberus ortizi</i> type 2	García San Nicolás, 1957	19,0	28,0	J/CO	8
L34		<i>Lauria cylindracea</i>	Da Costa, 1778	2,0	3,0	T/LC	2
L46		<i>Obba rota</i>	Broderip, 1841	24,0	30,0	F	2,5
L36		<i>Oestophora prietoi</i>	Ruiz, Arrebola & Puente, 2009	9,0	13,0	J	15
L70	3	<i>Otala ahmarica</i> kesebiana	Pallary, 1920		28,0	MC	3
C5	6	<i>Otala depageana</i>	Pallary, 1923	31,0	35,0	MC	3
C6		<i>Otala derenica</i> type 1	Pallary, 1920	26,0	33,0	MC	3
C14	6	<i>Otala derenica</i> type 2	Pallary, 1920	29,0	32,0	MC	6
C17		<i>Otala fauxnigra</i>	Chemnitz, 1786	25,0	38,0	MC	1
L84	3	<i>Otala fauxnigra</i> minor	Chemnitz, 1774	26,0	27,0	MC	4
C12		<i>Otala irreak</i> seir-repaired (several series)*		27,0	33,0	MC	2
C7		<i>Otala freydieri</i>	Pallary, 1923	25,0	31,0	MC	4
L9		<i>Otala lactea</i> *	Müller, 1774	21,0	39,0	J	0,5
C10		<i>Otala lactea murcica</i> black lip	Rossmässler, 1854	25,0	36,0	MU/MC	2
C1		<i>Otala lactea murcica</i> white lip	Rossmässler, 1854	25,0	38,0	MU	5
C2		<i>Otala lactea murcica</i> brown lip	Rossmässler, 1854	27,0	32,0	MU	3
L35	2	<i>Otala lucasii</i> white lip	Deshayes, 1850	24,0	28,0	MC	7
L123		<i>Otala punctata</i>	Müller, 1774	32,0	43,0	A/G	2
L81	1	<i>Otala riffensis</i>	Pallary, 1903		36,0	MC	4
L43	1	<i>Phaedusa lypra</i>	Mabille, 1887		22,0	CH	4
L71	1	<i>Phaedusa sheridani</i> shihi	Chang, 1982		15,0	TI	8
L42	1	<i>Plagiodontes dentatus</i> tesseirei	Marshal, 1930		15,0	UR	5
L31		<i>Pomatias elegans</i> W/O	Müller, 1774	11,0	19,0	CN	0,5
L75	2	<i>Pupinella swinhoi</i> W/O	Adams, 1866	9,0	10,0	TI	8
L30		<i>Rumina decollata</i>	Linné, 1758	27,0	40,0	J	0,5
L39		<i>Schistoloma alta</i> sibuyanica*	Bartsch, 1915	20,0	23,0	F	6
C13		<i>Sphincterochila baetica</i>	Rossmässler, 1835	20,0	26,0	AL	2
C15		<i>Sphincterochila candidissima</i> *	Draparnaud, 1801	19,0	23,0	J/G/AL	1,5
L88		<i>Theba pisana</i>	Müller, 1774	13,0	23,0	J / LC	0,5
L16		<i>Theba subdentata</i> helicella	Wood, 1828	18,0	20,0	AL	4
L48	2	<i>Tingitana</i> cf. <i>larochana</i>	Pallary, 1928	25,0	27,0	MC	6
L50		<i>Trochoidea elegans</i>	Draparnaud, 1801	7,0	12,0	M	1
L72		<i>Vallonia pulchella</i>	Müller, 1774	1,0	2,0	LC	1
L82	9	<i>Xerocrassa murcica</i>	Guirao in Schmidt, 1854	5,0	8,0	AL	4
L18		<i>Xerosecta arigonis</i>	Schmidt, 1853	16,0	24,0	J	2
L19		<i>Xerosecta promissa</i> white	Westerlund, 1893	11,0	15,0	G	1
L28		<i>Xerosecta reboudiana</i>	Bourguignat, 1863	9,0	11,0	G	2,5
L81	6	<i>Xerotricha apicina</i>	Lamarck, 1822	5,0	8,0	LC	1,5
L76		<i>Xerotricha conspurcata</i>	Draparnaud, 1801	3,0	6,0	J	1

Véase identificadores de localidades en la sección "Leyenda por países"

See place identifiers in the "Countries Caption" section